

ON AUTOMATIC CONTINUITY OF 3-HOMOMORPHISMS ON BANACH ALGEBRAS

JANKO BRAČIČ AND MOHAMMAD SAL MOSLEHIAN

ABSTRACT. A linear map $\varphi : \mathcal{A} \rightarrow \mathcal{B}$ between (Banach) algebras is called 3-homomorphism if $\varphi(abc) = \varphi(a)\varphi(b)\varphi(c)$ for each $a, b, c \in \mathcal{A}$. We investigate 3-homomorphisms on Banach algebras with bounded approximate identities and establish in two ways that every involution preserving homomorphism between C^* -algebras is norm decreasing.

1. INTRODUCTION.

In [3] the notion of 3-homomorphism between (Banach) algebras was introduced and some of their significant properties were investigated. A linear mapping $\varphi : \mathcal{A} \rightarrow \mathcal{B}$ between (Banach) algebras is called 3-homomorphism if $\varphi(abc) = \varphi(a)\varphi(b)\varphi(c)$ for each $a, b, c \in \mathcal{A}$. Obviously, each homomorphism is a 3-homomorphism. One can verify that if $\mathcal{A}$ is unital and φ is a 3-homomorphism then $\psi(a) = \varphi(1)\varphi(a)$ is a homomorphism.

There might exist 3-homomorphisms that are not homomorphisms. For instance, if $\psi : \mathcal{A} \rightarrow \mathcal{B}$ is a homomorphism, then $\varphi := -\psi$ is a 3-homomorphism which is not a homomorphism. As another example, assume that $\mathcal{A}$ and $\mathcal{B}$ are Banach algebras such that $\mathcal{A}^3 = \{0\}$ and $\mathcal{B}^3 = \{0\}$. Then each linear mapping between $\mathcal{A}$ and $\mathcal{B}$ is trivially a 3-homomorphism. One can find some non-trivial examples in [3].

In this paper, we investigate 3-homomorphisms on Banach algebras with bounded approximate identities and establish in two ways that every involution preserving homomorphism between C^* -algebras is norm decreasing.

Throughout the paper, we denote by T^* the dual of a linear mapping T and by $\mathcal{A}^*$ the dual of a Banach algebra $\mathcal{A}$. We also denote the involution on a C^* -algebra by $\#$.

2000 *Mathematics Subject Classification.* Primary 47B48, secondary 46L05, 16Wxx.

Key words and phrases. 3-homomorphism, homomorphism, continuity, Banach algebra, bounded approximate identity, second dual, Arens regularity, C^* -algebra.

2. 3-HOMOMORPHISMS ON C^* -ALGEBRAS

Suppose that φ is a non-zero 3-homomorphism from a unital algebra $\mathcal{A}$ to $\mathbb{C}$. Then $\varphi(1) = 1$ or -1 . Hence either φ or $-\varphi$ is a character on $\mathcal{A}$. If $\mathcal{A}$ is a Banach algebra, then φ is automatically continuous ([1, Theorem 2.1.29]). It may happen, however, that a 3-homomorphism is not continuous. For instance, let $\mathcal{A}$ be the algebra of all 3×3 matrices having 0 on and below the diagonal and $\mathcal{B}$ be the algebra of all $\mathcal{A}$ -valued continuous functions from $[0, 1]$ into $\mathcal{A}$ with sup norm. Then $\mathcal{B}$ is an infinite dimensional Banach algebra in which the product of any three elements is 0. Since $\mathcal{B}$ is infinite dimensional there are linear discontinuous maps and so discontinuous 3-homomorphisms from $\mathcal{B}$ into itself.

Using extreme points of the closed unit ball of a W^* -algebra $\mathcal{A}$ authors of [3] showed that if $\mathcal{B}$ is a C^* -algebra and $\varphi : \mathcal{A} \rightarrow \mathcal{B}$ is a weakly-norm continuous involution preserving 3-homomorphism, then $\|\varphi\| \leq 1$. Then they asked whether every preserving involution 3-homomorphism between C^* -algebras is continuous. We shall establish this by applying the strategy used by L. Harris in [2, Proposition 3.4] where it is shown that each J^* -homomorphism between J^* -algebras is norm decreasing.

Theorem 2.1. *Let $\varphi : \mathcal{A} \rightarrow \mathcal{B}$ be an involution preserving 3-homomorphism between unital C^* -algebras. Then $\|\varphi\| \leq 1$.*

Proof. Let $a \in \mathcal{A}$, let $\lambda > 0$ and let $\lambda \notin \sigma(a^\#a)$. Then $\lambda 1_{\mathcal{A}} - a^\#a$ has an inverse b . Then we have the following sequence of deductions:

$$\begin{aligned}
ab(\lambda 1_{\mathcal{A}} - a^\#a) &= a \\
(\lambda b - ba^\#a)a(\lambda b - ba^\#a) &= a \\
\varphi(\lambda b - ba^\#a)\varphi(a)(\lambda\varphi(b) - \varphi(b)\varphi(a^\#)\varphi(a)) &= \varphi(a) \\
\varphi(a^\#)(\varphi(\lambda b - ba^\#a)\varphi(a)\varphi(b))(\lambda 1_{\mathcal{B}} - \varphi(a^\#)\varphi(a)) &= \varphi(a^\#)\varphi(a) \\
\varphi(a^\#)\varphi(ab)(\lambda 1_{\mathcal{B}} - \varphi(a^\#)\varphi(a)) &= \varphi(a^\#)\varphi(a) \\
\lambda 1_{\mathcal{B}} - \varphi(a^\#)\varphi(ab)(\lambda 1_{\mathcal{B}} - \varphi(a^\#)\varphi(a)) &= \lambda 1_{\mathcal{B}} - \varphi(a^\#)\varphi(a) \\
1_{\mathcal{B}} &= \frac{1}{\lambda}(1_{\mathcal{B}} + \varphi(a^\#)\varphi(ab))(\lambda 1_{\mathcal{B}} - \varphi(a^\#)\varphi(a))
\end{aligned}$$

Since $\lambda 1_{\mathcal{B}} - \varphi(a^\#)\varphi(a)$ is self-adjoint, it follows that $\lambda 1_{\mathcal{B}} - \varphi(a)^\#\varphi(a)$ is invertible and so $\lambda \notin \sigma(\varphi(a)^\#\varphi(a))$. Hence

$$\sigma(\varphi(a)^\#\varphi(a)) \subseteq \sigma(a^\#a) \cup \{0\}.$$

Since the norm of a normal element c of a C^* -algebra is equal to its spectral radius $r(c)$ (see [1, Theorem 3.2.3]), we have

$$\|\varphi(a)\|^2 = \|\varphi(a^\#a)\| = r(\varphi(a)^\# \varphi(a)) \leq r(a^\#a) = \|a^\#a\| = \|a\|^2.$$

So φ is norm decreasing. $\square$

3. 3-HOMOMORPHISMS FROM ALGEBRAS WITH BOUNDED APPROXIMATE IDENTITY

Let $\mathcal{A}$ be a Banach algebra. Let $a \in \mathcal{A}$ and $\xi \in \mathcal{A}^*$ be arbitrary, then $a.\xi \in \mathcal{A}^*$ and $\xi.a \in \mathcal{A}^*$ are defined by $\langle a.\xi, x \rangle = \langle \xi, xa \rangle, x \in \mathcal{A}$ and $\langle \xi.a, x \rangle = \langle \xi, ax \rangle, x \in \mathcal{A}$.

For $\xi \in \mathcal{A}^*$ and $F \in \mathcal{A}^{**}$, functionals $F.\xi \in \mathcal{A}^*$ and $\xi.F \in \mathcal{A}^*$ are given by $\langle F.\xi, x \rangle = \langle F, \xi.x \rangle, x \in \mathcal{A}$ and $\langle \xi.F, x \rangle = \langle F, x.\xi \rangle, x \in \mathcal{A}$. Recall that the first Arens product of $F, G \in \mathcal{A}^{**}$ is $F \triangleleft G \in \mathcal{A}^{**}$, where

$$\langle F \triangleleft G, \xi \rangle = \langle F, G.\xi \rangle \quad (\xi \in \mathcal{A}^*).$$

Similarly, the second Arens product of $F, G \in \mathcal{A}^{**}$ is $F \triangleright G \in \mathcal{A}^{**}$, where

$$\langle F \triangleright G, \xi \rangle = \langle G, \xi.F \rangle \quad (\xi \in \mathcal{A}^*).$$

It is well known that $(\mathcal{A}^{**}, \triangleleft)$ and $(\mathcal{A}^{**}, \triangleright)$ are Banach algebras and that $\mathcal{A}$ is a closed subalgebra of each of them. The algebra $\mathcal{A}$ is Arens regular if the products $\triangleleft$ and $\triangleright$ coincide on $\mathcal{A}^{**}$.

A left approximate identity for $\mathcal{A}$ is a net $(e_\alpha) \subset \mathcal{A}$ such that $\lim_\alpha e_\alpha a = a$, for each $a \in \mathcal{A}$. If there exists $m > 0$ such that $\sup_\alpha \|e_\alpha\| \leq m$, the left approximate identity (e_α) is said to be bounded by m . The definitions of a right approximate identity and a bounded right approximate identity are similar. Recall from [1, Proposition 2.9.16] that $\mathcal{A}$ has a left approximate identity bounded by m if and only if $(\mathcal{A}^{**}, \triangleright)$ has a left identity of norm at most m . Similarly, $\mathcal{A}$ has a right approximate identity bounded by m if and only if $(\mathcal{A}^{**}, \triangleleft)$ has a right identity of norm at most m .

Lemma 3.1. *Let $\mathcal{A}$ and $\mathcal{B}$ be Banach algebras and $\varphi : \mathcal{A} \rightarrow \mathcal{B}$ be a 3-homomorphism. Then, for arbitrary $a, b \in \mathcal{A}$, $\xi \in \mathcal{B}^*$, and $F, G \in \mathcal{A}^{**}$, the following hold:*

- (i) $\varphi^*(\varphi(a)\varphi(b).\xi) = ab.\varphi^*(\xi)$;
- (ii) $\varphi^*(\xi.\varphi(a)\varphi(b)) = \varphi^*(\xi).ab$;
- (iii) $\varphi^*(\varphi(a).\xi.\varphi^{**}(F)) = a.\varphi^*(\xi).F$;
- (iv) $\varphi^*(\varphi^{**}(F).\xi.\varphi(a)) = F.\varphi^*(\xi).a$;

$$(v) \quad \varphi^*(\xi.\varphi^{**}(F) \triangleright \varphi^{**}(G)) = \varphi^*(\xi).F \triangleright G;$$

$$(vi) \quad \varphi^*(\varphi^{**}(F) \triangleleft \varphi^{**}(G).\xi) = F \triangleleft G.\varphi^*(\xi).$$

Proof. (i) Let $a, b \in \mathcal{A}$, and $\xi \in \mathcal{B}^*$ be arbitrary. Then, for each $x \in \mathcal{A}$,

$$\begin{aligned} \langle \varphi^*(\varphi(a)\varphi(b).\xi), x \rangle &= \langle \varphi(a)\varphi(b).\xi, \varphi(x) \rangle \\ &= \langle \xi, \varphi(x)\varphi(a)\varphi(b) \rangle \\ &= \langle \xi, \varphi(xab) \rangle \\ &= \langle \varphi^*(\xi), xab \rangle \\ &= \langle ab.\varphi^*(\xi), x \rangle. \end{aligned}$$

(iii) Let $a \in \mathcal{A}$, $\xi \in \mathcal{B}^*$, and $F \in \mathcal{A}^{**}$ be arbitrary. Then, for each $x \in \mathcal{A}$,

$$\begin{aligned} \langle \varphi^*(\varphi(a).\xi.\varphi^{**}(F)), x \rangle &= \langle \varphi(a).\xi.\varphi^{**}(F), \varphi(x) \rangle \\ &= \langle \varphi^{**}(F), \varphi(x)\varphi(a).\xi \rangle \\ &= \langle F, \varphi^*(\varphi(x)\varphi(a).\xi) \rangle \\ &= \langle F, xa.\varphi^*(\xi) \rangle \\ &= \langle a.\varphi^*(\xi).F, x \rangle. \end{aligned}$$

(v) Let $\xi \in \mathcal{B}^*$, and $F, G \in \mathcal{A}^{**}$ be arbitrary. Then, for each $x \in \mathcal{A}$,

$$\begin{aligned} \langle \varphi^*(\xi.\varphi^{**}(F) \triangleright \varphi^{**}(G)), x \rangle &= \langle (\xi.\varphi^{**}(F) \triangleright \varphi^{**}(G)), \varphi(x) \rangle \\ &= \langle \varphi^{**}(G), \varphi(x).\xi.\varphi^{**}(F) \rangle \\ &= \langle G, \varphi^*(\varphi(x).\xi.\varphi^{**}(F)) \rangle \\ &= \langle G, x.\varphi^*(\xi).F \rangle \\ &= \langle \varphi^*(\xi).F \triangleright G, x \rangle. \end{aligned}$$

The proofs of (ii), (iv), and (vi) are similar. □

Proposition 3.2. *Let $\mathcal{A}$ and $\mathcal{B}$ be Banach algebras. If $\varphi : \mathcal{A} \rightarrow \mathcal{B}$ is a 3-homomorphism, then φ^{**} is a 3-homomorphism from $(\mathcal{A}^{**}, \triangleleft)$ to $(\mathcal{B}^{**}, \triangleleft)$ and from $(\mathcal{A}^{**}, \triangleright)$ to $(\mathcal{B}^{**}, \triangleright)$.*

Proof. Let $F, G, H \in \mathcal{A}^{**}$ be arbitrary. Then, for each $\xi \in \mathcal{B}^*$,

$$\begin{aligned}
\langle \varphi^{**}(F \triangleleft G \triangleleft H), \xi \rangle &= \langle F \triangleleft G \triangleleft H, \varphi^*(\xi) \rangle \\
&= \langle F, G \triangleleft H, \varphi^*(\xi) \rangle \\
&= \langle F, \varphi^*(\varphi^{**}(G) \triangleleft \varphi^{**}(H), \xi) \rangle \\
&= \langle \varphi^{**}(F), \varphi^{**}(G) \triangleleft (\varphi^{**}(H), \xi) \rangle \\
&= \langle \varphi^{**}(F) \triangleleft \varphi^{**}(G), \varphi^{**}(H), \xi \rangle \\
&= \langle \varphi^{**}(F) \triangleleft \varphi^{**}(G) \triangleleft \varphi^{**}(H), \xi \rangle,
\end{aligned}$$

where the second equality holds by Lemma 5.1 (vi).

The proof of the second assertion is similar. $\square$

Theorem 3.3. *Let $\mathcal{A}$ and $\mathcal{B}$ be Banach algebras and $\varphi : \mathcal{A} \rightarrow \mathcal{B}$ be a 3-homomorphism. If $\mathcal{A}$ has a bounded left approximate identity, then*

$$\psi(F) := \varphi^{**}(L) \triangleright \varphi^{**}(F) \quad (F \in \mathcal{A}^{**})$$

*defines a homomorphism from $(\mathcal{A}^{**}, \triangleright)$ to $(\mathcal{B}^{**}, \triangleright)$, where L is a left identity in $\mathcal{A}^{**}$. Moreover,*

$$\varphi(a) = \varphi^{**}(L) \triangleright \psi(a) \quad (a \in \mathcal{A}).$$

Similarly, if $\mathcal{A}$ has a bounded right approximate identity, then

$$\psi(F) := \varphi^{**}(F) \triangleleft \varphi^{**}(R) \quad (F \in \mathcal{A}^{**})$$

*defines a homomorphism from $(\mathcal{A}^{**}, \triangleleft)$ to $(\mathcal{B}^{**}, \triangleleft)$, where R is a right identity in $\mathcal{A}^{**}$. Moreover,*

$$\varphi(a) = \psi(a) \triangleleft \varphi^{**}(R) \quad (a \in \mathcal{A}).$$

Proof. We shall prove only the first part of the theorem. Since φ is a 3-homomorphism the mapping φ^{**} is a 3-homomorphism from $(\mathcal{A}^{**}, \triangleright)$ to $(\mathcal{B}^{**}, \triangleright)$, by Proposition 3.2. By [1, Proposition 2.9.16], there exists a left identity L in $(\mathcal{A}^{**}, \triangleright)$. Thus

$$\psi(F) := \varphi^{**}(L) \triangleright \varphi^{**}(F) \quad (F \in \mathcal{A}^{**})$$

defines a homomorphism from $(\mathcal{A}^{**}, \triangleright)$ to $(\mathcal{B}^{**}, \triangleright)$ and

$$\varphi^{**}(F) = \varphi^{**}(L) \triangleright \psi(F) \quad (F \in \mathcal{A}^{**}).$$

Since the restriction of φ^{**} to $\mathcal{A}$ is φ , we have

$$\varphi(a) = \varphi^{**}(L) \triangleright \psi(a) \quad (a \in \mathcal{A}).$$

□

We now give another proof of Theorem 2.1:

Corollary 3.4. *Suppose that $\mathcal{A}$ and $\mathcal{B}$ are C^* -algebras. Then each involution preserving 3-homomorphism $\varphi : \mathcal{A} \rightarrow \mathcal{B}$ is bounded with norm $\|\varphi\| \leq 1$.*

Proof. Each C^* -algebra has approximate identity of bound 1 (cf. [1, Theorem 3.2.21]). By [1, Corollary 3.2.37], $\mathcal{A}$ and $\mathcal{B}$ are Arens regular and $\mathcal{A}^{**}$ and $\mathcal{B}^{**}$ are von Neumann algebras equipped with the involution that are defined in the following way (see [1, Page 349]). A linear involution on $\mathcal{A}^*$ is defined by

$$\langle \xi^\#, a \rangle = \overline{\langle \xi, a^\# \rangle} \quad (a \in \mathcal{A}, \xi \in \mathcal{A}^*),$$

and an involution on $\mathcal{A}^{**}$ is given by

$$\langle F^\#, \xi \rangle = \overline{\langle F, \xi^\# \rangle} \quad (F \in \mathcal{A}^{**}, \xi \in \mathcal{A}^*).$$

The involution on $\mathcal{B}^{**}$ is defined similarly. The algebra $\mathcal{A}^{**}$ is unital, with identity E of norm $\|E\| = 1$.

It is easy to check that φ^{**} is an involution preserving 3-homomorphism. Namely, for each $a \in \mathcal{A}$ and each $\xi \in \mathcal{B}^*$, we have

$$\langle \varphi^*(\xi^\#), a \rangle = \overline{\langle \xi, \varphi(a)^\# \rangle} = \overline{\langle \xi, \varphi(a^\#) \rangle} = \langle \varphi^*(\xi)^\#, a \rangle.$$

It follows, for each $F \in \mathcal{A}^{**}$ and each $\xi \in \mathcal{B}^*$, that

$$\langle \varphi^{**}(F^\#), \xi \rangle = \overline{\langle F, \varphi^*(\xi)^\# \rangle} = \overline{\langle F, \varphi^*(\xi^\#) \rangle} = \langle \varphi^{**}(F)^\#, \xi \rangle.$$

Since

$$\begin{aligned} \varphi^{**}(E)\varphi^{**}(F) &= \varphi^{**}(E)\varphi^{**}(FE^2) \\ &= \varphi^{**}(E)\varphi^{**}(F)\varphi^{**}(E)^2 \\ &= \varphi^{**}(EFE)\varphi^{**}(E) \\ &= \varphi^{**}(F)\varphi^{**}(E), \end{aligned}$$

for each $F \in \mathcal{A}^{**}$, we conclude that $\psi(F) = \varphi^{**}(E)\varphi^{**}(F)$ ($F \in \mathcal{A}^{**}$) defines an involution preserving homomorphism. Indeed, for each $F \in \mathcal{A}^{**}$, we have

$$\psi(F^\#) = \varphi^{**}(E)\varphi^{**}(F^\#) = (\varphi^{**}(F)\varphi^{**}(E)^\#)^\# = \psi(F)^\#.$$

By [1, Corollary 3.2.4], each involution preserving homomorphism between C^* -algebras is norm-decreasing. Thus

$$\|\varphi^{**}(F)\| = \|\varphi^{**}(E)\psi(F)\| \leq \|\varphi^{**}(E)\|\|F\| \quad (F \in \mathcal{A}^{**}).$$

Since

$$\psi(E) = \varphi^{**}(E)^2 = \varphi^{**}(E)^\# \varphi^{**}(E)$$

we have

$$1 \geq \|\psi(E)\| = \|\varphi^{**}(E)^\# \varphi^{**}(E)\| = \|\varphi^{**}(E)\|^2,$$

which gives $\|\varphi^{**}(E)\| \leq 1$. Thus,

$$\|\varphi^{**}(F)\| \leq \|\varphi^{**}(E)\|\|F\| \leq \|F\| \quad (F \in \mathcal{A}^{**})$$

□

Acknowledgement. The authors would like to thank the referees for their valuable comments and Professor John. D. Trout for his useful comment on Theorem 2.1.

REFERENCES

- [1] H. G. Dales, *Banach Algebras and Automatic Continuity*, London Math. Soc. Monographs 24, Clarendon Press, Oxford, 2000.
- [2] L. A. Harris, *A generalization of C^* -algebras*, Proc. Lond. Math. Soc. (3) **42** (1981), 331–361.
- [3] S. Hejazian, M. Mirzavaziri and M. S. Moslehian, *n -homomorphisms*, Bull. Iranian Math. Soc. **31** (2005), no. 1, 13–23.

DEPARTMENT OF MATHEMATICS, UNIV. OF LJUBLJANA, JADRANSKA UL. 19, 1111 LJUBLJANA, SLOVENIA

E-mail address: janko.bracic@fmf.uni-lj.si

DEPARTMENT OF MATHEMATICS, FERDOWSI UNIVERSITY, P. O. BOX 1159, MASHHAD 91775, IRAN;
CENTRE OF EXCELLENCE IN ANALYSIS ON ALGEBRAIC STRUCTURES (CEAAS), FERDOWSI UNIV., IRAN;
BANACH MATHEMATICAL RESEARCH GROUP (BMRG), MASHHAD, IRAN.

E-mail address: moslehian@ferdowsi.um.ac.ir